

# MEETING AGENDA

The General Managers Advisory Committee Meeting on a Dog Park will be held at the **893 Southwood Boulevard (IVGID Admin Building)** starting at 4:00 p.m. on Wednesday, November 16, 2022.

A. Roll call of the committee members

Committee Members: Leighton Pratt, Simi Balter, Myles Riner, Judith Miller, Michelle Lintzner, Janet Pahl, Indra Winquest, IVGID District General Manager, and Shelia Leijon, Director of Parks and Recreation.  
Administrative Support Staff: Susan Herron

Rules of Engagement: All committee members shall be respectful of one another and treat each other with kindness. No one member shall talk over another member. All ideas are welcome however, they may not be discussed in depth or in detail depending upon time and/or merit and that determination is solely held with the General Manager. All committee members are equal participants.

B. Committee discussion on Diamond Peak site

C. Discussion with Nevada Department of Wildlife representative Samuel Sedillo regarding stream and lake health

D. Set next meeting date/time – November 30, 2022 (proposed)

E. Adjournment

X Fencing along the existing parking lot

Existing Parking

Large Dog Park

Small Dog Park

Possible Deck and Restroom location



# Dog Wastes and Water Quality: *Evaluating the Connection at Lake Tahoe*

**Mark Walker**, Water Quality Specialist, University of Nevada Cooperative Extension, Department of Natural Resources and Environmental Sciences, University of Nevada, Reno, NV

**Lynell Garfield**, Graduate Student, Environmental Sciences Graduate Program, Department of Natural Resources and Environmental Sciences, University of Nevada, Reno, NV

*This fact sheet discusses research that examined potential water contamination from a popular dog exercise area. The study consisted of water sampling and estimating the total amount of dog wastes deposited over the course the study, which lasted 14 months. The study found that approximately 100.1 lbs (45.5 kg) of dog wastes (as dry matter) accumulated and that accumulations were highly localized. However, it found no link between bacteria in water and waste accumulations. Populations of *E. coli*, a microbe that indicates water contamination from the feces of mammals, degrade quickly when evaporative conditions are high. Land managers can control waste accumulations to minimize the potential for water pollution using several strategies, which are briefly reviewed.*

## DOG WASTES AND WATER QUALITY

People have become concerned about the effects of accumulated dog wastes on water quality. Dog wastes, like any wastes from mammals, may contain very large numbers of microbes, some of which can cause disease. If enough dog waste accumulates in an area, they could contaminate water, which could affect human health. Examples of diseases that can be transmitted from dogs to humans through feces include salmonellosis, giardiasis and cryptosporidiosis (see [www.agnr.umd.edu/CES/Pubs/PDF/FS703.pdf](http://www.agnr.umd.edu/CES/Pubs/PDF/FS703.pdf) for a list of the disease-causing microorganisms that could be present in dog wastes).

Open spaces such as parks and walkways are often used as exercise areas for dogs. In many places, dog-owners are encouraged to collect and dispose of dog wastes, with judiciously placed bag dispensers and waste cans. However, some areas do not have such facilities.

Our study investigated the potential connection between accumulations of dog wastes and the quality of water in a small tributary to Lake Tahoe (figure 1), by estimating the amount of dog wastes that accumulated in the area

and testing water for *E. coli*.

The results have some interesting implications for how bag stations and receptacles should be sited to help dog owners keep trails and recreation areas clean.

## HOW AND WHERE THE STUDY WAS DONE

We conducted the study in an area criss crossed by a popular trail system managed by the U.S. Forest Service at Lake Ta-

hroe. The trails led to the lake and passed by a small pond in the middle of the stream course of Burke Creek (Figure 2). Before conducting the study we observed that many residents drove their dogs to a small parking area to walk, run and cycle with them. Because there were no bag dispensers or waste cans, very few people




Figure 1: Location of the study site on east side of Lake Tahoe, in Nevada.




Figure 2: The Burke Creek Recreational Area study site at Lake Tahoe. Triangles represent plots where dog wastes were collected. This figure shows the estimated amounts of dog wastes accumulated during a two week period ending on 6/21/05.

collected and disposed of wastes. In fact, when walking the trails, we found heavy accumulations of dog wastes in many parts of the study area.

The study area covered 8.8 acres (Figure 2). Within the boundaries of the study area, we selected 15 places to collect dog wastes and five sites to sample water. The water sampling sites started from where Burke Creek entered the study area (called *Below Highway (BH)* in Figures 2 and 3), to where it left the study area (*Below Bridge (BB)*). Every two weeks we collected wastes from the sampling points and took water. We also determined how many *E. coli* could be found in dog wastes. Finally, we carried out experiments to determine how long *E. coli* could survive in canine feces when exposed to different rates of evaporation.

We estimated the total accumulations of wastes by developing maps of the mass of wastes per unit area. We reported results of water sample analysis as “colony forming units (CFU)/100 ml of water.” A colony forming unit is a microbial colony—a shiny round spot—that forms on the surface of the filters used to filter up to 100 ml of water (slightly less than seven tablespoons). The filters retain *E. coli* and are incubated with nutrients to promote rapid growth, so a CFU represents at least one *E. coli* cell that was able to reproduce. As an example, if a water sample has 10 CFU/100 ml, this indicates that there were at least 10 *E. coli* cells that formed colonies when a water sample was filtered.

### THE RESULTS

**Waste Accumulations:** Dog wastes were distributed very unevenly throughout the study site. Most wastes accumulated in areas that were either very close to trailheads or where trails crossed (Figure 2). This is likely because of canine territory marking behavior and preference for certain kinds of toilet areas.




Figure 3: Accumulations of dog waste (left y-axis) and water sampling results at the inlet to the study site (BH) and outlet from the study site (BB) over the 14 month study period (x-axis).

Overall, approximately 100 pounds of wastes, as dry matter, accumulated in the study area over the course of 14 months. The accumulations differed by season, with much less in winter months than in other seasons. This was likely because the site was covered with snow and inaccessible.

As shown in Figure 2, wastes did not accumulate evenly across the entire 8.8 acre site. Accumulations were limited to areas such as trail junctions, areas near parking lots and entry points to the recreation area. The areas of highest accumulation varied between sampling periods, but the total area in which wastes were found was always much smaller than the 8.8 acre site.

***E. coli* In Wastes and Water Content:** Dog wastes were an average of 47% water. A gram (slightly less than 0.04 ounces) of fresh feces contained an average of 50 million CFU/gram with a range of two million to 200 million CFU/g. The wide range can be attributed to the highly variable nature of dog food, digestive health and diets.

***Link Between Water Samples and Waste Accumulations:*** Sampling results did not show water contamination or a link between accumulated dog wastes and *E. coli* in Burke Creek (Figure 3). In fact, we found that water leaving the study area had fewer CFU/100 ml than water entering the study area. This may have been due to a wetland through which the creek meandered and a small pond in the stream course that was designed to trap sediment. Although numbers of *E. coli* CFU/100 ml were occasionally high, no single sample from water leaving the study area exceeded federal guidelines set to prevent illness from contact with water (a geometric average of 126 CFU/100 ml).

***E. coli* Survival In Dog Waste:** We found nearly complete loss of *E. coli* in dog feces within 60 hrs of exposure to evaporation rates of 0.08 inches/day and within 15 hours for 0.30 inches/day. Average daily evaporation rates at Lake Tahoe range from approximately 0.20 in/day to close to zero during wet cool months typical

of late fall and spring months. Although temperature had a small effect on the rate of *E. coli* die-off, water content had the biggest effect on how quickly *E. coli* disappeared from feces.

## **WHAT THE RESULTS MEAN**

The results suggest that under the right circumstances *E. coli* in dog wastes may die quickly as moisture evaporates from feces, with complete die-off even before feces are completely desiccated. This helps to explain why we found no link between *E. coli* in water samples and the accumulations of dog wastes in the study area drained by Burke Creek.

In order for water to be contaminated, bacteria must enter the water. Soils at the site were sandy and, unless frozen, very unlikely to have water running off them. At the site, commonly used toilet areas may have exposed feces to high evaporation rates that quickly killed *E. coli*, especially during the hot, dry months of summer, when the most feces accumulated.

The study has several limitations that are important to understand. First, the survival studies took place under carefully controlled conditions in a laboratory. This was necessary to be sure that death of *E. coli* could be linked with evaporation rates. Second, the samples used in the study were smaller than an average dog feces. This means that the effects of evaporation on *E. coli* survival were likely to have been exaggerated. Third, although *E. coli* is considered an indicator of contamination by feces, it survives environmental stresses differently than disease-causing microorganisms. In fact, microorganisms such as *Cryptosporidium* and *Giardia* survive environmental stresses at much more extreme levels and for longer durations than those that were used for these experiments.

The results do not mean that dog wastes cannot contaminate water. In fact, under the right circumstances, such as the cool, moist periods in spring and late fall, feces could enter water supplies before pathogenic mi-

croorganisms can be killed by desiccation or other stresses.



### **STRATEGIES FOR REDUCING THE RISK OF CONTAMINATION WITH DISEASE CAUSING MICROORGANISMS**

The risk of contamination could be high when evaporation rates are low and the potential for runoff from snowmelt or rainfall is high, for example during fall and spring months. The risk could be reduced significantly, using several strategies either singly or in combination.

**First**, waste accumulates in specific areas—for example at trail junctions or near trail heads. These areas are ideal places to install bag dispensing stations and litter bins.

**Second**, it is possible to manipulate where wastes accumulate in certain areas. For example, setting a pole surrounded by a light coating of sand on the soil encourages dogs to defecate near the pole, prompted by marking behavior (see *Animal Waste Collection* under “Additional Resources”). If sited away from streams and rills on the landscape and accompanied by bag dispensing stations and waste collection bins, such sites could ensure that wastes accumulate away from water bodies and are removed. Also, dogs prefer to defecate in areas where grass is longer (about 4”) and not mowed frequently. Maintaining a toilet area away from streams and paved surfaces and providing owners with the

means to collect and remove wastes can substantially reduce the potential for contamination.

**Third**, a public awareness campaign can encourage dog owners to collect and properly dispose of wastes. This could involve working closely with land management agencies (local municipal and county parks departments). One of the most important parts of starting a campaign is to be sure that resources are available to sustain the program, including maintaining bagging stations and disposing of wastes.

---

### **For further information please contact:**

**Mark Walker**, Associate Professor, State Cooperative Water Quality Extension Specialist, Department of Natural Resources and Environmental Science, University of Nevada, Reno  
Phone: (775) 784-1938  
Email: [mwalker@cabnr.unr.edu](mailto:mwalker@cabnr.unr.edu)

### **Additional Resources:**

Public Open Space and Dogs, 1995. Harlock Jackson Pty Ltd Ch. 7: Specific Guidelines for the selection, design and management of individual parks. <http://www.petnet.com.au/openspace/frontis.html>  
Pollution Prevention: Animal Waste Collection [www.stormwatercenter.net/Pollution\\_Prevention\\_Factsheets/AnimalWasteCollection.htm](http://www.stormwatercenter.net/Pollution_Prevention_Factsheets/AnimalWasteCollection.htm) (last accessed 4/2008)

**This publication produced with support from:** *The Nevada Agricultural Experiment Station at the*



*University of Nevada and a Regional Water Quality Coordination Grant provided by the U.S. Department of Agriculture’s Cooperative State Research, Education and Extension Service.*

---

*The University of Nevada, Reno is an Equal Opportunity/Affirmative Action employer and does not discriminate on the basis of race, color, religion, sex, age, creed, national origin, veteran status, physical or mental disability, and in accordance with University policy, sexual orientation, in any program or activity it operates. The University of Nevada employs only United States citizens and aliens lawfully authorized to work in the United States.*

*Copyright © 2008, University of Nevada Cooperative Extension*